

High frequency monitoring of carbon (iv) oxide evolution from a tropical soil amended with organomineral fertilizer in a screenhouse Ile-Ife, Nigeria

Tijani, F. O.*, Okunkenu, A. S., Afolabi, F., Akinde, B. P. and Oyedele, D. J.

Department of Soil Science and Land Resources Management, Obafemi Awolowo University, Ile-Ife, Osun State, Nigeria.

*Corresponding author: Email: dftijani@oauife.edu.ng

Copyright © 2024 Tijani et al. This article remains permanently open access under the terms of the [Creative Commons Attribution License 4.0](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received 5th July 2024; Accepted 10th August 2024

ABSTRACT: The study compared responses of carbon (iv) oxide evolved from an Ultisol sampled at different time scales and the soil properties when amended with organomineral fertilizer (OMF); thus tracking the carbon footprint of the amendment in ambient air. The carbon (iv) oxide efflux was measured titrimetrically in a static chamber set up in a screenhouse at sub-daily, daily and weekly time scales. At the same time, the antecedent soil properties were determined using standard methods. The treatments involved OMF addition at three rates (0 kg OMF, 40 kg Urea-N + 2.5 tons/ha organic fertilizer and 40 kg Urea-N + 5 tons/ha organic fertilizer) based on some Nigerian indigenous vegetable nutrient requirements. The result showed that a more detailed and accurate understanding of CO₂ evolution processes was revealed at daily sampling resolution while sub-daily variation occurred in response to sub-daily variation in ambient temperature. Organomineral fertilizer at 40 kg/ha Urea + 2.5 tons/ha OF rate was comparatively safer in C management through the reduction in the CO₂ released thus constituting a better alternative in terms of the greenhouse effect. Thus, the study recommends at least daily monitoring for a detailed understanding of CO₂ evolution dynamics and management and identifies OMF rate with less CO₂ release as a comparatively better alternative in the production of those vegetables in the environmental context.

Keywords: Carbon (iv) oxide release, environment, organomineral fertilizers, sampling frequency, soil.

Abbreviations: OF, Organic fertilizer, OMF, Organomineral fertilizer; CO₂, Carbon (iv) Oxide, SOM, Soil organic matter, GHG, Greenhouse gas

INTRODUCTION

To meet the need for food by the global teeming population, agriculture has relied on the use of mineral fertilizers to sustain crop output and this is not without environmental implications (Rahman and Zhang, 2018; Ilahi *et al.*, 2020). One of such impacts is global warming. The reality, importance and impact of global warming in the world cannot be ignored. Greenhouse gases, which include carbon dioxide (CO₂), methane (CH₄) and nitrous oxides are major contributors to greenhouse gas emissions (GHGs) (Shi *et al.*, 2006; Bréchet *et al.*, 2021;

Mohammed *et al.*, 2021). Notably, CO₂ efflux from soil to the atmosphere is a major component of GHGs, and it is a crucial pathway of the carbon (C) cycle (Shi *et al.*, 2006; Liu *et al.*, 2011; Mohammed *et al.*, 2021). The soil CO₂ efflux that gets to the atmosphere is produced by heterotrophic (decomposer organisms) and autotrophic (living roots and mycorrhizae) activities which depend on the substrate quality and the environmental conditions Subke *et al.*, 2003; Kim *et al.*, 2010; Lopez-Canfin *et al.*, 2018). It makes up about 20-40% of the total annual input

of CO₂ into the atmosphere (Subke *et al.*, 2003; Kim *et al.*, 2010; Carlson *et al.*, 2016). Lorenz and Lal (2014) reported that CO₂ contributes about 60% to greenhouse gas emissions (GHG) and that any activity that results in C sequestration could be an effective mitigation strategy for global climate change. Soil temperature and moisture are among the most important factors controlling soil CO₂ efflux (Berglund *et al.*, 2010; Oyedele and Tijani, 2010; Gomes *et al.*, 2016; Yang *et al.*, 2017). Previous studies indicated that soil temperature is the major environmental driving factor influencing CO₂ efflux (Shi *et al.*, 2006; Hursh *et al.*, 2016; Kuzyakov *et al.*, 2019; Gołasa *et al.*, 2021). A number of organic materials could be incorporated into the soil to increase C sequestration in agricultural lands. This in turn reduces the release of CO₂ into the atmosphere (Fageria, 2012; Gayathri *et al.*, 2021). Such materials include biochar, animal manure, plant residues etc. as against inorganic fertilizers.

To increase crop yield, it is becoming standard practice to apply excessive amounts of synthetic fertilizer. Only one or two nutritional components can be supplied to the crop by employing chemical fertilizers alone (Hou *et al.*, 2010), thus leading to nutrient imbalance (Pahalvi *et al.*, 2021). Their use has been found to be effective as a short-term solution and it demands consistent use on a long-term basis (Makinde *et al.*, 2007; Olowoake *et al.*, 2015; Ghosh *et al.*, 2021). However, they are most times too expensive for low-income, small-scale farmers (Makinde *et al.*, 2007; Ghosh *et al.*, 2021). Conversely, organic amendments are considerably less costly and supply significant carbon and vital nutrients to meet the needs of soil microbes (Ilahi *et al.*, 2020). Their use cannot be overemphasized because of their ability to improve the physical, chemical and biological properties of the soil (Wang *et al.*, 2019; Zhang *et al.*, 2019). They improve the conditions in which crops grow and develop which in turn culminates into better production of economic plants (Olowoake, 2014; Ulzen *et al.*, 2020). For example, organic fertilizer application (and the resultant C input) increases soil organic carbon (SOC) pool, whereas, the inorganic fertilizer application may increase, decrease, or have no significant effect on SOC stock (Olowoake, 2014; Yang *et al.*, 2017). Zougmore *et al.* (2009) reported that organic fertilizer application improves water-holding capacity (WHC) and infiltration as a result of well-developed aggregates. The drawback of a low supply of plant nutrients has been reported for organic manures (Mahmood *et al.*, 2017). However, a nutrient management technique that employs the integration of fertilizers (both organic and inorganic) so as to safeguard the sustainable condition of the soil and give plants the ideal amount of nutrients needed to sustain yield productivity seems promising (Ayinla *et al.*, 2019; Panta and Parajulee, 2021). Thus, the complementary use of organic and inorganic fertilizer (organomineral) generally improves the yield and quality of crops and increases SOC stock as compared with the sole application of mineral

fertilizer (Olowoake, 2014; Oshunsanya and Akinrinola, 2014; Yang *et al.*, 2017). Babalola *et al.* (2007), Hou *et al.* (2010), Gai *et al.* (2018) and Smith *et al.* (2020) reported that SOC retention, soil C sequestration and soil infiltration are significantly affected by organomineral fertilizer (OMF). Also, vegetable growers reported that organomineral fertilizers (OMF) improved physico-chemical characteristics of soil and the growth of vegetables (*Amaranthus cruentus*, *Abelmoschus esculentus*) compared to the control (non-additive/untreated soil) (Makinde *et al.*, 2011; Olowoake, 2014; Olowoake *et al.*, 2015).

One of the crops gaining prominent attention in Nigeria is vegetables. Vegetables are an important part of healthy eating and have been found to be a source of many nutrients including potassium, fibre, folate and vitamins A, E and C (Olowo *et al.*, 2022). Nutrients in vegetables can help to maintain healthy blood pressure, reduce blood cholesterol levels and lower the risk of heart disease (Moyo *et al.*, 2020). In Nigeria, there are two broad categories of vegetables, the exotic and the indigenous. Exotic vegetables that are not indigenous to the country (Sayeed *et al.*, 2021). Some have been domesticated and grown in some parts of the country (such as broccoli, kale, strawberries etc.) while some are imported (such as apples, Egyptian grapes, etc.) (Amfo and Baba-Ali, 2020). On the other hand, the indigenous vegetables are native to Nigeria (Adenegan *et al.*, 2016). Generally, these indigenous vegetables are cheap to procure and excellent sources of proteins, carbohydrates, minerals, vitamins and dietary fibres, and sources of income for smallholder farmers (Amujoyegbe *et al.*, 2015; Lawal *et al.*, 2018). Some examples of prominent Nigerian indigenous vegetables include *Amunututu* or Ceylon/Malabar spinach (*Basella alba*), *Ila* or Okra (*Abelmoschus esculentus* L.), *Efinrin* or African Basil (*Ocimum gratissimum* L.), *Igba* or Garden egg (*Solanum macrocarpon*), *Odu* or Nightshade (*Solanum nigrum*), *Ewuro* or Bitter leaf (*Vernonia amygdalina*), *Ugwu* or Fluted pumpkin (*Telfaria occidentalis*), *Ebolo* or Ragleaf (*Crassocephalum crepidioides*), *Isapa* or Roselle (*Hibiscus sabdariffa*), *Eggplant leaf* (*Solanum melongena*), *Wild leaf Marugbo* or *Eweta* (*Clerodendrum volubile*), *Sokoyokoto* or Cocks' comb (*Celosia argentea* L.), *Gbure* or Water leaf (*Talinum triangulare*), *Tete* or Amaranth (*Amaranthus hybridus*), *Snake tomato* or *Tomato elejo* (*Trichosanthes cucumerina*) (Lawal *et al.*, 2018).

Studies on these indigenous vegetables have shown that the application of OMF allows a comprehensive soil fertility management strategy which will give benefits of applying an organic fertilizer as well as applying a little dose of inorganic fertilizer (Oyedele *et al.*, 2017). According to a study on fertilizer micro-dosing of indigenous Nigerian vegetables by Adebooye *et al.* (2018), it was observed that applying Urea (inorganic fertilizer) at 20 kg, 40 kg and 60 kg Urea-N/ha, along with basal application of organic fertilizer, gave better income

(\$40.51, \$45.89 and \$61.08, respectively, in a plot size of 6 m²) than 80 kg Urea-N/ha (\$39.52 from 6 m² plot size). This result confirmed that fertilizer micro-dosing can be used to sustainably produce these indigenous vegetables, particularly when co-applied with organic manure (Akponikpe *et al.*, 2018; Ayanwale *et al.*, 2018). Organic fertilizers have been reported to improve carbon sequestration (Wang *et al.*, 2019; Ghosh *et al.*, 2021); thereby, mitigating the release of CO₂ to the atmosphere. Despite the promising results obtained from OMF application for vegetable production, little research has been done on the carbon sequestration potential of the OMF, particularly on CO₂ efflux. With differences in organic materials and their combinations with mineral fertilizers such as OMF, and possible variation in carbon loss over time, there is the need for more investigation not only on their capacity to improve soil properties and yield but also on the implications of these amendments on the carbon in the atmosphere. Therefore, the knowledge of CO₂ emissions from soil and how this emission changes with time are of great importance to estimate atmospheric CO₂ concentration and ascertain carbon sequestration of the capacity of the soil system (Shi *et al.*, 2006; Kuzyakov *et al.*, 2019). In most soil and water studies, researchers have relied on low-frequency monitoring such as monthly or yearly (Bowes *et al.*, 2009; Bah *et al.*, 2012; Sale *et al.*, 2021). However, such low-frequency data can obscure a timely response and decisions that pertain to rapid changes in soil CO₂ efflux (Bah *et al.*, 2012; Cândido *et al.*, 2020). Soil carbon dioxide efflux which is an index of microbial activities and nutrient mineralization, is closely associated with soil functioning, and carbon sequestration, and has environmental implications (Gomes *et al.*, 2016; Lopez-Canfin *et al.*, 2018). The possible obscurity that exists in low-frequency or coarse monitoring of CO₂ efflux has suggested that carbon efflux processes occur at very fine resolution and require high-frequency monitoring (Chen *et al.*, 2019). High-frequency monitoring is necessary for a better understanding of nutrient dynamics and carbon response in the soil as well as its release into the atmosphere (Zeng *et al.*, 2016; Sith *et al.*, 2017; Bréchet *et al.*, 2021). Therefore, this study hypothesized that temporal variability at the sub-daily scale exists with CO₂ efflux and that its loss to the atmosphere varies with the rate of fertilizer application. The objectives of this study were to (1) investigate the difference in sub-daily, daily and weekly monitoring of CO₂ and (2) assess the temporal variability of CO₂ efflux as affected by temperature and organomineral fertilizers, and their effect on the selected properties of soil.

MATERIALS AND METHODS

Study site and soil

The study was conducted in a screenhouse of the Faculty

of Agriculture, Obafemi Awolowo University, Ile-Ife, Nigeria (latitude 7°31'14.76" and longitude 4°31'49.13"). The maximum daily temperatures ranged between 27 and 35°C, and the minimum temperature between 18.9 and 23.3°C. The soil used for the experiment was an Ultisol, a low base-status forest soil developed from coarse-grained gneiss and granite parent rocks (Okusami and Oyediran, 1985). It is classified as Typic Paleustult (Soil Survey Staff, 1975) and also classified as the Iwo series (Okusami and Oyediran, 1985).

Sample collection and preparation, experimental design and layout

Bulk soil samples were taken from a field that has been subjected to continuous cultivation (with maize) for over 15 years (degraded) in the Teaching and Research Farm (T & R), OAU, Ile-Ife, Nigeria. The sampling was done on a relatively flat terrain to a depth of 15 cm. The samples were air-dried, gently crushed and then passed through a 2-mm mesh sieve. Composite samples were taken from the bulked soil samples for physical and chemical analyses.

Urea fertilizer and un-amended compost (organic fertilizer (OF); Sunshine Grade B) developed by the Ondo State Government of Nigeria and were used to prepare the OMF. The results of analyses of the organic fertilizer are Nitrogen (3.5%), Phosphorous (1.0%) and Potassium (2.5%) (Ondo State Government, 2012). Table 1 shows the treatment combinations used for the experiment. Each treatment was thoroughly mixed with 4 kg air-dried soil and repacked in plastic pots of 4.5 m³ volume (perforated at the bottom for free drainage and air movement). The experiment was arranged in a completely randomized design (CRD) consisting of three treatments replicated four times giving a total of 12 pots. They were maintained at 75% field capacity (FC).

Measurement of released CO₂ during soil incubation

The amount of CO₂ released during soil incubation was measured as described by Oyedele and Tijani (2010). Twelve potted samples of non-amended soil and soils amended with organomineral fertilizer that were maintained at 75% field capacity (FC) were placed in a static chamber. During the CO₂ sampling (incubation) period, specifically 2 hours, the pots were kept in the air-tight condition. Each pot had a beaker containing 10 ml of 0.5 M NaOH solution. An empty pot (without soil) that had a beaker (10 ml of 0.5 NaOH solution) was taken as blank (Kim *et al.*, 2010; Oyedele and Tijani, 2010; Gelfand *et al.*, 2015). The ambient temperature was assessed by measuring the temperature of water in a beaker within the chamber using a mercury thermometer. The incubation was carried out at a sub-daily time scale (7.00 GMT and

Table 1. The description of the study's treatment.

No.	Treatment	Code
1	Control (0 kg Urea-N/ha + 0 tons/ ha OF)	OMF1
2	First Rate (40 kg Urea-N/ha + 2.5 tons/ha OF)	OMF2
3	Second Rate (40 kg Urea-N/ha + 5 tons/ha OF)	OMF3

12.00 GMT) and each incubation lasted 2 hours for each measurement for thirty-one days for the determination of CO₂ evolution. The emitted CO₂ from the soil was trapped in the base solution in the beaker, in each static chamber which was later titrated with 0.5 M H₂SO₄ using methyl red indicator.

Laboratory analyses

Preliminary soil sampling was carried out on the soil to determine its antecedent soil properties (such as soil texture, bulk density, soil organic matter, pH, and available phosphorus) before imposing the treatment. The soil pH was determined by a digital pH meter (Walk lab Ti 9000) in a 2:1 0.01 M CaCl₂ solution to soil suspension calibrated to buffers pH 4 and 7 (Thomas, 1996). The other soil parameters were; particle size distribution determined using a modified hydrometer method as described by Gee and Or (2002); soil organic matter determined using the dichromate oxidation method (Nelson and Sommers, 1996) and available phosphorus determined by Bray-1 method (Bray and Kurtz, 1945; Kuo, 1996). The FC was the volumetric moisture content after gravitational drainage as adapted from Flint and Flint (2002). The bulk density was obtained via the use of the core method described by Stolte and Veerman (1997).

Statistical analysis

The data obtained was subjected to analysis of variance (ANOVA). Means were separated using Duncan's Multiple Range Test (DMRT) at $p \leq 0.05$.

RESULTS AND DISCUSSION

The section revealed that significant differences in CO₂ release from soil to the environment, that was hidden when sampling at weekly and other coarser sampling frequencies were noticeable at a finer resolution such as daily and sub-daily monitoring scales. Furthermore, the loss of the gas increased at noon in response to the increase in temperature and also soil treated with OMF2 (40 kg urea + 2.5 tons organic fertilizer) was environmentally preferable because the soil released less carbon than OMF3 (40 kg urea + 5 tons organic fertilizer).

Although the study was conducted in a screenhouse and it will also be interesting to make observations on the field, this monitoring scale is not often considered in soil studies and it is the area this study draws its strength. Details of the findings are presented in the sub-sections below.

Properties of the soil before imposing the treatments

The soil properties before application of the amendments are presented in Table 2 indicating a sandy loam soil having a bulk density of 1.56 gcm⁻³, which was close to the critical value (1.63 gcm⁻³) when soil is considered impaired for good crop performance. Soil organic matter (SOM) and available P were below the critical level reported by Adepetu *et al.* (2014). More detailed discussion of these antecedent soil properties and further investigation on them, which include the soil texture, bulk density, soil hydraulic conductivity, pH, SOM and available P have been reported (Tijani *et al.*, 2024).

Carbon emission at a relatively low sampling frequency in response to the application of organomineral fertilizer

Table 3 reveals the CO₂ loss at the beginning and end of the experiment. Though week 5 has a lower soil CO₂ loss than that of week 1 for all the treatments and there was no significant difference ($p > 0.05$) among the treatments at week 5, there was less percentage loss of C (73.8% loss) between week 1 and week 5 for soil on OMF2 as against OMF1 and OMF3 which have 84.7% and 89.3% losses of C, respectively, at the same temperature. Changes in C release to the atmosphere from the treated soils and its dynamics could also be due to the fact that SOM was quickly used up at the beginning by microorganisms for their metabolic activities which resulted in increased respiration, which only became more pronounced as the rate increased, as observed in OMF3. Further, at week 1, OMF3 emitted more CO₂ than both OMF1 and OMF2 while the two were not significantly different. This indicates that OMF2 posed a relatively less C threat to the environment probably due to the lower rate of the fertilizer applied, relative to OMF3 and the slow nutrient release attribute of the organic component of the fertilizer. The latter may have caused the difference between OMF2 and OMF1 in per

Table 2. The antecedent properties of soil.

Soil property	Value
Sand (gkg ⁻¹)	702
Clay (gkg ⁻¹)	197
Silt (gkg ⁻¹)	101
Textural class	Sandy loam
Bulk density (Mgm ⁻¹)	1.56
SOM (%)	1.55
pH	5.1
Avail. P (Bray-1) (mg/kg)	8.3

SOM = soil organic matter, Avail. P = Available phosphorous.

Table 3. Soil emission of CO₂ and ambient temperature at the beginning and end of the study.

Treatments	Week 1	Week 5
Temp. (°C)	31	29
CO ₂ (kg ha day ⁻¹)		
OMF1	255.0b	39.0a
OMF2	240.4b	62.8a
OMF3	345.3a	37.1a

OMF1, OMF2 and OMF3 = (0 tons/ha OF + 0 kg Urea-N/ha), (2.5 tons/ha OF + 40 kg Urea-N/ha) and (5 tons/ha OF + 40 kg Urea-N/ha). Means in a column with the same alphabet are not statistically different at $p \leq 0.05$.

Table 4. Evolution of CO₂ and associated ambient temperature from the soil at weekly time scale.

Treatments	Week 1	Week 2	Week 3	Week 4	Week 5
Temp. (°C)	31	29	31	30	29
CO ₂ (kg ha day ⁻¹)					
OMF1	255.0b	88.7a	118.2a	59.1a	39.0a
OMF2	240.4b	85.0a	113.1a	77.6a	62.8a
OMF3	345.3a	62.83a	110.5a	70.2a	37.1a

OMF1, OMF2 and OMF3 = (0 tons/ha OF + 0 kg Urea-N/ha), (2.5 tons/ha OF + 40 kg Urea-N/ha) and (5 tons/ha OF + 40 kg Urea-N/ha). Means in a column with the same alphabet are not statistically different at $p \leq 0.05$.

cent C loss between weeks 1 and 5. The reduction in CO₂ evolution from 255.0, 240.4 and 345.3 kg ha⁻¹ day⁻¹ in week 1 to 39.0, 62.8 and 37.1 kg ha⁻¹ day⁻¹ in week 5 for OMF1, OMF2 and OMF3, respectively, is in agreement with Ghosh *et al.* (2021) who linked such occurrence to SOM dynamics in soil.

Table 4 reveals the weekly CO₂ evolution and associated ambient temperature. There was no significant difference ($p > 0.05$) in CO₂ evolution between the treatments across the weeks of sampling except week 1 (Table 4). Furthermore, there was an initial increase in CO₂ efflux in week 1 that was higher than those in weeks 2 to 5. This may be due to the rapid disintegration of OMF by

microorganisms that needed nitrogen for their activities, such that there was an increase in microbial activity leading to the release of more CO₂. Pandey *et al.* (2010) and Zamanian *et al.* (2018) reported that the addition of inorganic fertilizer especially nitrogen appears to have a priming effect on increasing CO₂ efflux from the soil as it releases its nutrients faster within a short time. CO₂ evolution responded to mild temperature changes from week 2 to week 4. This aligned with the reports that ascertained that a strong positive correlation exists between soil temperature and CO₂ efflux (Fenn *et al.*, 2010; Oyedele and Tijani, 2010; Galic *et al.*, 2020; Bréchet *et al.*, 2021).




Figure 1 (A & B). Effect of varied rate of organomineral fertilizer on soil CO₂ efflux among the treatments at sub-daily time scale. A – morning sampling; B – afternoon sampling; OMF1 - Control, OMF2 – organomineral fertilizer at 40 kg Urea-N/ha + 2.5 ton/ha OF, OMF3 – organomineral fertilizer at 40 kg Urea-N/ha + 5 ton/ha OF.

Soil CO₂ efflux of the treated soil at daily and sub-daily time scales

Figure 1 shows the trend in CO₂ loss to the ambient air in the morning and at noon at a daily time scale in response to the different rates of the OMF treatment. The comparison of the CO₂ signals between the morning and noon represents the sub-daily monitoring frequency. At either morning or noon, days in which there were significant ($p < 0.05$) emissions of CO₂ from the soil to the ambient air and potentially to the atmosphere were about 10% of the experimental period. The emission was dominated by OMF3. As the temperature rose at noon, carbon loss from soil more than doubled. The occurrence of higher carbon efflux was predominant in the early days of the experiment and this was more pronounced at noon. This is to re-emphasize the strong relationship that exists between temperature and CO₂ evolution (Subke *et al.*, 2003; Shi *et al.*, 2006; Saiz *et al.*, 2007; Kim *et al.*, 2010;

Bréchet *et al.*, 2021) and the importance of proper sampling regime for the all-encompassing understanding of CO₂ dynamics. Figure 2A to C depict the variation in CO₂ signals from the soil when sampled in the morning and noon (sub-daily monitoring of CO₂) for each of the treatments. The CO₂ evolution in the first three days (at the beginning) at noon was significantly higher ($p < 0.05$) than in the morning (with lesser temperatures) for all treatments (a few days after imposing OMF) implying an initial rise in CO₂ evolution for all the treatments. This indicates the effect of microbial decomposition on the SOM. Considering the number of days (10 days; 32.3% of the experimental period) that soil CO₂ efflux was significantly higher at noon than CO₂ efflux in the morning for each of OMF1 (the control) and OMF2 against OMF3 that has 13 days (41.9% of the experimental duration) of significant CO₂ efflux, therefore, OMF3 created more CO₂ concern to the environment. Consequently, OMF2 (2.5 tons/ ha + 40 kg Urea-N/ha) added relatively no additional carbon




Figure 2 (A, B & C). Sub-daily sampling of soil CO₂ efflux from soil treated with organomineral fertilizer. A – Control, B – organomineral fertilizer at 40 kg Urea-N/ha + 2.5 ton/ha OF, C – organomineral fertilizer at 40 kg Urea-N/ha + 5 ton/ha OF.

burden to the environment (due to its rate), while at the same time improving soil properties (Tijani *et al.*, 2024), implying it could lower the amount of the greenhouse gas and its effect to the environment. These higher significant differences recorded (at noon) showed the variation that exists at different times of the day which is a pointer to the higher potential threat of the evolution of CO₂ to the environment in warmer climates. However, CO₂ release

from soil treated with OMF3 translated to the highest cumulative carbon mineralization during the incubation period. This also indicates that the mixture of organic and mineral fertilizers applied could improve soil microbial activity (Smith *et al.*, 2020), although with attendant higher CO₂ loss, unlike the comparatively lower mineralization that was observed in soils having OMF2 (compared to OMF3).




Figure 3 (A, B & C). Comparison of CO₂ efflux of the treatments at daily and weekly time scale. A – Control, B – organomineral fertilizer at 40 kg Urea-N ha⁻¹ + 2.5 ton ha⁻¹ OF, C - organomineral fertilizer at 40 kg Urea-N ha⁻¹ + 5 ton ha⁻¹ OF.

This indicates that OMF2 had the greatest potential in safer C management than OMF3. The resultant effect is a reduction in the amount of CO₂ evolution and a potential reduction in leached nutrients released to the environment. Therefore, management practices such as balanced application of fertilizer could play important roles in soil C sequestration or safe C management and thereby GHG mitigation, and reduction in the pollution of ground/surface water (Ghosh *et al.*, 2010; Fageria, 2012; Lorenz and Lal, 2014). The results also revealed that temporal variability exists with soil CO₂ efflux at sub-daily scale. However, OMF2 exhibited comparatively less variations between morning and noon, than OMF3 across the days of the experiment, despite the higher temperature recorded at noon. This could be due to less quantity of SOM added

through OMF2 compared to that of OMF3 leading to a reduction in the amount of CO₂ released to the environment. Zhang *et al.* (2014), Wang *et al.* (2016) and Liang *et al.* (2022) also reported that low fertilizer application could lower the release of GHGs and reduce global warming.

Comparisons between measurements of CO₂ evolution at daily and weekly time scales

The daily sampling clearly shows more details of the trend and decline of CO₂ evolution as against the weekly sampling (Figure 3A, B and C) which obscures such details. The combined low and high sampling frequency of CO₂

efflux revealed that high-frequency monitoring was more accurate (or revealing) in the understanding of carbon dynamics. Although the weekly monitoring reasonably captured the trend in carbon dynamics, there was an underestimation of carbon (CO₂) emissions from the soil. This underscores the importance of this monitoring approach in carbon management in soil as done for rivers by Kooistra *et al.* (2012) and Grift *et al.* (2015) who advocated for combined low and high-frequency monitoring for crop status in precision agriculture and estimates of nutrients in lowland rivers. Therefore, fine resolutions (such as sub-daily or daily measurements) would facilitate the synthesis of relevant information (Blackford *et al.*, 2017). This will endeavor users to make informed agronomic and economic decisions for better carbon and nutrient management, and profitable agricultural production and ecosystem health (Bah *et al.*, 2012). It is, therefore, reasonable to state that such high-resolution monitoring would be required for soil CO₂ dynamics variability (Tang *et al.*, 2003; Cao *et al.*, 2021). Sub-daily soil CO₂ efflux monitoring has also been suggested by other authors Chédin *et al.* (2003), Cao *et al.* (2021) and Chamizo *et al.* (2022).

Conclusions

This study elucidates the effect of OMF on soil CO₂ efflux (C dynamics). It was observed that OMF2 (40 kg Urea-N/ha + 2.5 tons of OF) was at a safer rate in terms of CO₂ evolution than OMF3 (40 kg Urea-N/ha + 5 tons/ha of OF). High-resolution monitoring (e.g. daily sampling) of CO₂ efflux was more revealing and accurate than low resolution monitoring and it is necessary for holistic understanding and management of carbon exchange between the soil and the atmosphere. Therefore, a cautious management strategy in the use of OMF could play a vital role in GHG mitigation in view of the fertilizer rate for the production of Nigerian indigenous vegetables in the environmental context.

CONFLICT OF INTEREST

The authors declared that no competing interest exists.

REFERENCES

- Adebooye, O. C., Akponikpe, P. I., Aluko, E. R., Oyedele, D., & Peak, D. J. (2018). CIFSrf final technical report: Scaling up fertilizer micro-dosing and indigenous vegetable production and utilization in West Africa (CIFSrf Phase 2). Retrieved from <https://idl-bnc-idrc.dspacedirect.org/server/api/core/bitstreams/a8138aa7-37aa-4374-811c-578d0543fa4f/content>
- Adenegan, K. O., Adegbite, O. O., & Ibe, R. B. (2016, August). Demand for selected exotic fruits and vegetables among government workers: a quadratic almost ideal demand system approach. In *III All Africa Horticultural Congress 1225* (pp. 399-406).
- Adepetu, J. A., Adetunji, M. T., & Ige, D. V. (2014). *Soil fertility and crop nutrition*. Jumak Publisher, Ibadan, Nigeria. p. 560.
- Akponikpe, P. I., Oyedele, D., Adebooye, O. C., & Peak, D. J. (2018). Final report of agronomic trial and water management of fertilizer micro-dosing technology on traditional leafy vegetables. Retrieved from <https://idl-bnc-idrc.dspacedirect.org/server/api/core/bitstreams/94fe454a-d5c3-48ae-8085-35b060b92f13/content>.
- Amfo, B., & Baba Ali, E. (2021). Technology adoption by indigenous and exotic vegetable farmers. *International Journal of Vegetable Science*, 27(2), 105-119.
- Amujoyegbe, B. J., Oyedele, D. J., Idowu, M. K., Ayinde, J. O., & Adebooye, O. C. (2015). On-farm adoption of under-utilized indigenous vegetable production among small holder farmers in Nigeria: Implication for economic empowerment and genetic conservation. *Journal of Agricultural Extension and Rural Development*, 7(9), 283-289.
- Ayanwale, A., Ajekiigbe, N., Oyedele, D., & Adebooye, C. (2018). Economic impacts of fertilizer micro-dosing in the production of underutilized indigenous vegetables in south west Nigeria: An ex-ante approach. In *III All Africa Horticultural Congress 1225* (pp. 303-308).
- Ayinla, A., Alagbe, I. A., Olayinka, B. U., Lawal, A. R., Aboyeji, O. O., & Etejere. (2018). Effects of organic, inorganic and organo-mineral fertilizer on the growth, yield and nutrient composition of *Corchorus Olitorious* (L.). *Ceylon Journal of Science*, 47(1), 13-19.
- Babalola, O., Oshunsanya, S. O., & Are, K. (2007) Effects of vetiver grass (*Vetiveria nigritana*) strips, vetiver grass mulch and an organomineral fertilizer on soil, water and nutrient losses and maize (*Zea mays*, L) yields, *Soil & Tillage Research*, 96, 6-18.
- Bah, A., Balasundram, S. K. & Husni, M. H. A. (2012) Sensor Technologies for Precision Soil Nutrient Management and Monitoring, *American Journal of Agricultural and Biological Sciences*, 7(1), 43-49.
- Berglund, Ö., Berglund, K., & Klemetsson, L. (2010). A lysimeter study on the effect of temperature on CO₂ emission from cultivated peat soils. *Geoderma*, 154(3-4), 211-218.
- Blackford, J., Artioli, Y., Clark, J., & de Mora, L. (2017). Monitoring of offshore geological carbon storage integrity: Implications of natural variability in the marine system and the assessment of anomaly detection criteria. *International Journal of Greenhouse Gas Control*, 64, 99-112.
- Bowes, M. J., Smith, J. T., & Neal, C. (2009). The value of high-resolution nutrient monitoring: A case study of the River Frome, Dorset, UK. *Journal of Hydrology*, 378, 82-96.
- Bray, R. H., & Kurtz, L. T. (1945) Determination of total, organic, and available forms of phosphorus in soils. *Soil Science Journal*, 59, 39-45.
- Bréchet, L. M., Daniel, W., Stahl, C., Burban, B., Goret, J., Salomón, R. L., & Janssens, I. A. (2021). Simultaneous tree stem and soil greenhouse gas (CO₂, CH₄, N₂O) flux measurements: a novel design for continuous monitoring towards improving flux estimates and temporal resolution. *New Phytologist*, 230(6), 2487-2500.
- Cândido, B. M., Quinton, J. N., James, M. R., Silva, M. L. N., de Carvalho, T. S., de Lima, W., & Eltner, A. (2020). High-resolution monitoring of diffuse (sheet or interrill) erosion using

- structure-from-motion. *Geoderma*, 375, 114477.
- Cao, M., Lei, J., He, Q., Zeng, Z., Lü, X., & Jiang, Y. (2021). Rainfall-driven and hydrologically-controlled variations in cave CO₂ sources and dynamics: Evidence from monitoring soil CO₂, stream flow and cave CO₂. *Journal of Hydrology*, 595, 126060.
- Carlson, K. M., Gerber, J. S., Mueller, N. D., Herrero, M., MacDonald, G. K., Brauman, K. A. & West, P. C. (2016). Greenhouse gas emissions intensity of global croplands. *Nature Climate Change*, 7(1), 63-68.
- Chamizo, S., Rodríguez-Caballero, E., Sánchez-Cañete, E. P., Domingo, F., & Cantón, Y. (2022). Temporal dynamics of dryland soil CO₂ efflux using high-frequency measurements: Patterns and dominant drivers among biocrust types, vegetation and bare soil. *Geoderma*, 405, 115404.
- Chédin, A., Saunders, R., Hollingsworth, A., Scott, N., Matricardi, M., Etcheto, J., Clerbaux, C., Armante, R., & Crevoisier, C. (2003). The feasibility of monitoring CO₂ from high-resolution infrared sounders. *Journal of Geophysical Research: Atmospheres*, 108(D2), 4064.
- Chen, Y., Guerschman, J. P., Cheng, Z., & Guo, L. (2019). Remote sensing for vegetation monitoring in carbon capture storage regions: A review. *Applied Energy*, 240, 312-326.
- Fageria, N. K. (2012). Role of soil organic matter in maintaining sustainability of cropping systems. *Communications in Soil Science and Plant Analysis*, 43(16), 2063-2113.
- Fenn, K. M., Malhi, Y., & Morecroft, M. D. (2010). Soil CO₂ efflux in a temperate deciduous forest: Environmental drivers and component contributions. *Soil Biology and Biochemistry*, 42(10), 1685-1693.
- Gai, X., Liu, H., Liu, J., Zhai, L., Yang, B., Wu, S., & Wang, H. (2018). Long-term benefits of combining chemical fertilizer and manure applications on crop yields and soil carbon and nitrogen stocks in North China Plain. *Agricultural Water Management*, 208, 384-392.
- Galic, M., Mesic, M., & Zgorelec, Z. (2020). Influence of organic and mineral fertilization on soil greenhouse gas emissions. A review. *Agriculturae Conspectus Scientificus*, 85(1), 1-8.
- Gayathri, R., Mahboob, S., Govindarajan, M., Al-Ghanim, K. A., Ahmed, Z., Al-Mulhm, N. & Vijayalakshmi, S. (2021). A review on biological carbon sequestration: A sustainable solution for a cleaner air environment, less pollution and lower health risks. *Journal of King Saud University-Science*, 33(2), 101282.
- Gee, G. W., & Or, D. (2002). Particle size analysis. In: Dane, J., & Topp, G. C. (eds.). *Methods of Soil Analysis. Book Series: 5. Part 4* (pp. 255-293). Soil Science Society of America, USA.
- Gelfand, I., Cui, M., Tang, J. & Robertson, G.P. (2015). Short term drought response of N₂O and CO₂ emissions from mesic agricultural soils in the US Midwest, *Agriculture, Ecosystems and Environments*, 212, 127-133.
- Ghosh, M., Ashiq, W., Bhogilal Vasava, H., Gamage, D. N. V., Patra, P. K., & Biswas, A. (2021). Short-Term carbon sequestration and changes of soil organic carbon pools in rice under integrated nutrient management in India. *Agriculture*, 11(4), 348.
- Ghosh, S., Wilson, B. R., Mandal, B., Ghoshal, S. K., & Grown, I. (2010) Changes in soil organic carbon pool in three long-term fertility experiments with different cropping systems and inorganic and organic soil amendments in the eastern cereal belt of India, *Australian Journal of Soil Research*, 48, 413-420.
- Gołasa, P., Wysokiński, M., Biełkowska-Gołasa, W., Gradziuk, P., Golonko, M., Gradziuk, B. & Gromada, A. (2021). Sources of greenhouse gas emissions in agriculture, with particular emphasis on emissions from energy used. *Energies*, 14(13), 3784.
- Gomes, L. de C., Cardoso, I. M., Mendonça, E. de S., Fernandes, R. B. A., Lopes, V. S., & Oliveira, T. S. (2016). Trees modify the dynamics of soil CO₂ efflux in coffee agroforestry systems. *Agricultural and Forest Meteorology*, 224, 30-39.
- Griff, B. V., Broers, H. P., Berendrecht, W. L., Rozemeijer, J. C., Osté, L. A., & Griffioen, J. (2015) High-frequency monitoring reveals nutrient sources and transport processes in an agriculture-dominated lowland water system, *Hydrology Earth System Sciences*, 12, 8337-8380.
- Hou, X., Han, X., Li, H., & Xing, B. (2010) Composition and Organic Carbon Distribution of Organomineral Complex in Black Soil under Different Land Uses and Management Systems, *Communications in Soil Science and Plant Analysis*, 41, 1129-1143.
- Hursh, A., Ballantyne, A., Cooper, L., Maneta, M., Kimball, J., & Watts, J. (2016). The sensitivity of soil respiration to soil temperature, moisture, and carbon supply at the global scale. *Global Change Biology*, 23(5), 2090-2103.
- Ilahi, H., Hidayat, K., Adnan, M., Rehman, F., Tahir, R., Saeed, M. S., Shah, S. W. A., & Toor, M. D. (2020). Accentuating the impact of inorganic and organic fertilizers on agriculture crop production: A review. *Indian Journal of Pure and Applied Bioscience* 9(1), 36-45.
- Kim, D. G., Mu, S., Kang, S., & Lee, D. (2010). Factors controlling soil CO₂ effluxes and the effects of rewetting on effluxes in adjacent deciduous, coniferous, and mixed forests in Korea. *Soil Biology and Biochemistry*, 42(4), 576-585.
- Kooistra, L., Beza, E. A., Verbesselt, J., van den Borne, J., & van der Velde, W. (2012). Integrating remote-, close range-and in-situ sensing for high-frequency observation of crop status to support precision agriculture. In *Proceedings Sensing a Changing World, Wageningen, The Netherlands, 9-11 May, 2012* (pp. 15-20).
- Kuo, S. (1996). Phosphorus determination. In: Bartels J. M (ed.). *Method of soil analysis. Chemical methods, Part 3* (pp. 869-919). Soil Science Society of America, USA.
- Kuzyakov, Y., Horwath, W. R., Dorodnikov, M., & Blagodatskaya, E. (2019). Review and synthesis of the effects of elevated atmospheric CO₂ on soil processes: No changes in pools, but increased fluxes and accelerated cycles. *Soil Biology and Biochemistry*, 128, 66-78.
- Lawal, O. M., Idowu-Adebayo, F., & Enujiugha, V. N. (2018). Nutritional assessment of Nigerian ethnic vegetable soups (Marugbo, Tete and Ila). *Journal of Nutrition, Food and Lipid Science*, 1(32-39), 10-33513.
- Liang, Z., Jin, X., Zhai, P., Zhao, Y., Cai, J., Li, S., Yang, S., Li, C., & Li, C. (2022) Combination of organic fertilizer and slow-release fertilizer increases pineapple yields, agronomic efficiency and reduces greenhouse gas emissions under reduced fertilization conditions in tropical areas. *Journal of Cleaner Production*, 343, 131054.
- Liu, J., Jiang, P., Wang, H., Zhou, G., Wu, J., Yang, F. & Qian, X. (2011). Seasonal soil CO₂ efflux dynamics after land use change from a natural forest to Moso bamboo plantations in subtropical China. *Forest Ecology and Management*, 262, 1132-1137.
- Lopez-Canfin, C., Sánchez-Cañete, E. P., Serrano-Ortiz, P., López-Ballesteros, A., Domingo, F., Kowalski, A. S., & Oyonarte, C. (2018). From microhabitat to ecosystem: Identifi-

- fying the biophysical factors controlling soil CO₂ dynamics in a karst shrubland. *European Journal of Soil Science*, 69(6), 1018-1029.
- Lorenz, K., & Lal, R. (2014). Biochar application to soil for climate change mitigation by soil organic carbon sequestration. *Journal of Plant Nutrition and Soil Science*, 177(5), 651-670.
- Mahmood, F., Khan, I., Ashraf, U., Shahzad, T., Hussain, S., Shahid, M., Abid, M., & Ullah, S. (2017). Effects of organic and inorganic manures on maize and their residual impact on soil physico-chemical properties. *Journal of soil science and plant Nutrition*, 17(1), 22-32.
- Makinde, E. A., Ayeni, L. S., & Ojeniyi, S. O. (2011). Effects of organic, organomineral and NPK fertilizer treatment on the nutrient uptake of *Amaranthus cruentus* (L) on two soil types in Lagos, Nigeria. *Journal of Central European Agriculture*, 12(1), 114-123.
- Makinde, E. A., Ayoola, O. T., & Akande, M. O. (2007). Effects of organomineral fertilizer application on the growth and yield of 'egusi' melon. *Australian Journal of Basic and Applied Sciences*, 1(1), 15-19.
- Mohammed, S., Gill, A. R., Alsafadi, K., Hijazi, O., Yadav, K. K., Hasan, M. A., & Harsanyi, E. (2021). An overview of greenhouse gases emissions in Hungary. *Journal of Cleaner Production*, 314, 127865.
- Moyo, S. M., Serem, J. C., Bester, M. J., Mavumengwana, V. & Kayitesi, E. (2020). African green leafy vegetables health benefits beyond nutrition. *Food Reviews International*, 37(6), 601-618.
- Nelson, D. W., & Sommers, L. E. (1996). Total carbon, organic carbon and organic matter. In: Sparks, D. L. (ed.). *Methods of soil analysis. Book Series: 5. Part 3* (2nd ed) (pp. 961-1010). Madison (WI): American Society of Agronomy and Soil Science Society of American, USA.
- Okusami, T. A., & Oyediran, G. O. (1985). Slope-soil relationships on an Aberrant Toposequence in Ife area of South Western Nigeria: Variabilities in soil properties. *Ife Journal of Agriculture*, 7(1), 1-15.
- Olowo, S. F., Omotayo, A. O., Lawal, I. O., & Aremu, A. O. (2022). Improving rural livelihood through the cultivation of indigenous fruits and vegetables: evidence from Ondo State, Nigeria. *Agriculture*, 12(3), 372.
- Olowoake, A. A. (2014). Influence of organic, mineral and organomineral fertilizers on growth, yield, and soil properties in grain amaranth (*Amaranthus cruentus*. L), *Journal of Organics*, 1(1), 39-47.
- Olowoake, A. A., Ojo, J. A., & Osunlola, O. S. (2015). Growth and yield of okra (*Abelmoschus esculentus* L.) as influenced by NPK, jatropha cake and organomineral fertiliser on an Alfisol in Ilorin, Southern Guinea Savanna of Nigeria. *Journal of Organic Systems*, 10(1), 3-8.
- Ondo State Government (2012). Waste to wealth. Ondo State Waste Management Bulletin on the use of Organic Fertilizer and Organomineral Fertilizer, Ondo State, Nigeria, p. 10.
- Oshunsanya, S. O., & Akinrinola, T. B. (2014). Changes in soil physical properties under yam production on a degraded soil amended with organomineral fertilizer. *Global Journal of Agricultural Research and Reviews*, 2(1), 87-93.
- Oyedele, D. J., & Tijani, F. O. (2010). Soil carbon dioxide evolution in response to agricultural land use, in: Adebooye, O. C., Taiwo, K. A., & Fatufe, A. A. (eds.). *Biotechnology development and threat of climatic change in Africa: The case of Nigeria*. Cuvillier Verlag Gottingen, 2, 211-215.
- Oyedele, D. J., Idowu, M. K., Tijani, F. O., & Egbebi, I. (2017). Design and assessment of a capillary irrigation system for fertilizer micro-dosing in vegetable production. *ASA, CSSA and SSSA International Annual*.
- Pahalvi, H. N., Rafiya, L., Rashid, S., Nisar, B., & Kamili, A. N. (2021). Chemical fertilizers and their impact on soil health. *Microbiota and Biofertilizers, Vol 2: Ecofriendly tools for reclamation of degraded soil environs*. Pp. 1-20.
- Pandey, R. R., Sharma, G., Singh, T. B. & Tripathi, S. K. (2010). Factors influencing soil CO₂ efflux in a northeastern Indian oak forest and plantation, *African Journal of Plant Science*, 4(8), 280-289.
- Panta, S., & Parajulee, D. (2021). Integrated Nutrient Management (INM) in soil and sustainable agriculture. *International Journal of Applied Science and Biotechnology*, 9(3), 160-165.
- Rahman, K. A., & Zhang, D. (2018). Effects of fertilizer broadcasting on the excessive use of inorganic fertilizers and environmental sustainability. *Sustainability*, 10(3), 759.
- Saiz, G., Black, K., Reidy, B., Lopez, S., & Farrell, E. P. (2007). Assessment of soil CO₂ efflux and its components using a process-based model in a young temperate forest site. *Geoderma*, 139(1-2), 79-89.
- Sale, T., Gallo, S., Askarani, K. K., Irianni-Renno, M., Lyverse, M., Hopkins, H., ... & Burge, S. (2021). Real-time soil and groundwater monitoring via spatial and temporal resolution of biogeochemical potentials. *Journal of Hazardous Materials*, 408, 124403.
- Sayeed, A., Islam, M. S., Uddin, M. N., Proshad, R., Kundu, S., Islam, M. N., & Hossen, M. M. (2021). Nutritional status of exotic and indigenous vegetables. *International Journal of Vegetable Science*, 27(1), 86-95.
- Shi, P. L., Zhang, X. Z., Zhong, Z. M., & Ouyang, H. (2006). Diurnal and seasonal variability of soil CO₂ efflux in a cropland ecosystem on the Tibetan Plateau. *Agricultural and Forest Meteorology*, 137(3-4), 220-233.
- Sith, R., Yamamoto, T., Watanabe, A., Nakamura, T., & Nadaoka, K. (2017). Analysis of red soil sediment yield in a small agricultural watershed in Ishigaki Island, Japan, using long-term and high resolution monitoring data. *Environmental Processes*, 4(2), 333-354.
- Smith, W. B., Wilson, M., & Pagliari, P. (2020). Organomineral fertilizers and their application to field crops. *Animal manure: Production, Characteristics, Environmental Concerns, and Management*, 67, 229-243.
- Soil Survey Staff (1975). *Soil taxonomy. A basic system for making and interpreting soil surveys*. Soil Conservation Service, Handbook 436, United States Department of Agriculture, Washington DC.
- Stolte, J., & Veerman, G. (1997). *Manual of soil physical measurements*, Version 3. Wageningen, D. L.O. Staring Centre. *Technical Document*, 37.
- Subke, J. A., Reichstein, M., & Tenhunen, J. D. (2003). Explaining temporal variation in soil CO₂ efflux in a mature spruce forest in Southern Germany. *Soil Biology and Biochemistry*, 35(11), 1467-1483.
- Tang, J., Baldocchi, D. D., Qi, Y., & Xu, L. (2003). Assessing soil CO₂ efflux using continuous measurements of CO₂ profiles in soils with small solid-state sensors. *Agricultural and Forest Meteorology*, 118(3-4), 207-220.
- Thomas, G. W. (1996). Soil pH and soil acidity. In: Sparks, D. L.

- (ed.). *Methods of soil analysis: Chemical methods, part 3*. Soil Science Society of America, Madison WI., USA.
- Tijani, F. O., Okunkenu, A. S., Afolabi, F., Akinde, B. P., & Oyedele D. J. (2024). Evaluating unsaturated hydraulic conductivity of a tropical soil amended with organomineral fertilizer at various time scales in a screenhouse in Ile-Ife, Nigeria. *Ife Journal of Science*, 26(2), 521-535.
- Ulzen, J., Abaidoo, R. C., Ewusi-Mensah, N., & Masso, C. (2020). Combined application of inoculant, phosphorus and organic manure improves grain yield of cowpea. *Archives of Agronomy and Soil Science*, 66(10), 1358-1372.
- Wang, B., Wan, Y., Qin, X., Gao, Q., Liu, S., & Li, J. (2016). Modifying nitrogen fertilizer practices can reduce greenhouse gas emissions from a Chinese double rice cropping system. *Agriculture, Ecosystems & Environment*, 215, 100-109.
- Wang, H., Xu, J., Liu, X., Zhang, D., Li, L., Li, W., & Sheng, L. (2019). Effects of long-term application of organic fertilizer on improving organic matter content and retarding acidity in red soil from China. *Soil and Tillage Research*, 195, 104382.
- Wang, N., Li, J. Y., & Xu, R. K. (2009). Use of agricultural by-products to study the pH effects in an acid tea garden soil. *Soil Use and Management*, 25(2), 128-132.
- Yang, M., Li, Y., Li Y., Cheng, S. X., Yue, T., Fu, W., Jiang, P., & Zhou, G. (2017). Effects of inorganic and organic fertilizers on soil CO₂ efflux and labile organic carbon pools in an intensively managed Moso bamboo (*Phyllostachys pubescens*) plantation in subtropical China. *Communications in Soil Science and Plant Analysis*, 48(3), 332-344.
- Zamanian, K., Zarebanadkouki, M., & Kuzyakov, Y. (2018). Nitrogen fertilization raises CO₂ efflux from inorganic carbon: A global assessment. *Global Change Biology*, 24(7), 2810–2817.
- Zeng, C., Liu, Z., Zhao, M., & Yang, R. (2016). Hydrologically-driven variations in the karst-related carbon sink fluxes: Insights from high-resolution monitoring of three karst catchments in Southwest China. *Journal of Hydrology*, 533, 74-90.
- Zhang, J., Bei, S., Li, B., Zhang, J., Christie, P., & Li, X. (2019). Organic fertilizer, but not heavy liming, enhances banana biomass, increases soil organic carbon and modifies soil microbiota. *Applied Soil Ecology*, 136, 67-79.
- Zougmore, R., Mando, A., & Stroosnyder, L. (2009) Soil nutrient and sediment loss as affected by erosions barriers and nutrients source in semi-arid Burkina Faso. *Arid Land Research and Management*, 23, 85-101.