

Influence of poultry manure rates on postharvest quality and shelf-life of green pepper (*Capsicum annuum*) fruits

Evans Ntim Amedor^{1*}, John Yao Afetsu², Robert Akayim Awasina³, Salifu Abdallah⁴ and Austin Christopher Gyamfi⁵

^{1,5}Department of Science, Bia Lamplighter College of Education, Sefwi Debiso, Ghana.

²Department of Science, St. Teresa's College of Education, Hohoe, Ghana.

^{3,4}Department of Science, Gambaga College of Education, Gambaga, Ghana.

*Corresponding author: Email: evamedor999@gmail.com

Copyright © 2025 Amedor et al. This article remains permanently open access under the terms of the [Creative Commons Attribution License 4.0](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

Received 6th August 2024; Accepted 15th January 2025

ABSTRACT: The research investigated the impact of different poultry manure rates on the post-harvest quality and shelf-life of green pepper (*Capsicum annuum*) fruits. The experiment utilised a 3 × 1 factorial design, implemented within a Randomized Complete Block Design (RCBD) framework. The study included three replications to ensure statistical validity. The treatment details comprised: T₁ (0 kg/ha, control), T₂ (1.2 kg/ha poultry manure), and T₃ (2.4 kg/ha poultry manure) with a common green pepper variety, California Wonder. Results indicated that poultry manure application significantly influenced postharvest parameters. The 2.4 kg/ha poultry manure treatment exhibited the lowest weight loss and maintained the longest shelf-life and marketable quality after 12 days of storage. Additionally, fruits from this treatment demonstrated the highest vitamin C and crude fibre contents. In contrast, the 1.2 kg/ha poultry manure treatment resulted in the highest chlorophyll content. The results of this study indicate that the application of poultry manure at a rate of 2.4 kg/ha yields the most favourable outcomes for green pepper fruits. This specific rate appears to enhance post-harvest quality characteristics and prolong the shelf-life of the peppers.

Keywords: Chlorophyll, crude fibre, physiological weight loss, treatments, vitamin C.

INTRODUCTION

Green pepper (*Capsicum annuum*), also known as bell pepper, is valued for its nutritional value and culinary adaptability (Brezeanu *et al.*, 2022). Green peppers are a nutritional powerhouse containing a diverse array of essentials including minerals (potassium, calcium, phosphorus, and iron), vitamins (A, B, and C), and carbs, proteins, and lipids (Danfulloh *et al.*, 2022). As the most prevalent antioxidant, vitamin C is recognized to help prevent cancer and other diseases linked to weakened immune systems in humans (Anaya-Esparza *et al.*, 2021; Jamshidovich, 2023). The quality and storability of green pepper fruits are significantly impacted by pre-harvest management practices, which may involve the use of organic fertilizers like poultry manure.

Poultry manure is a valuable organic fertilizer that

increases crop productivity and soil fertility since it is high in potassium, phosphorus, and nitrogen (Ngongolo and Mrimi, 2024; Cairo-Cairo *et al.*, 2023). In addition to improving soil structure and water-holding capacity, its application influences the biochemical composition and physiological responses of plants. This results in improved water uptake and decreased water stress in plants, which is reflected in firmer and more resilient fruits after harvest (Arshad *et al.*, 2024). The rate at which green pepper plants are able to absorb nutrients from the soil is directly impacted by the amount of poultry manure applied to the soil. Nutrient deficits brought on by insufficient application rates of poultry manure may hinder plant growth and fruit development (Akinfasoye, 2019). Conversely, high manure application rates can cause nutritional imbalances,

contaminate the environment, and lower fruit quality because of nutrient toxicity or physiological issues (Risse *et al.*, 2006, Drózdź, 2020). In a study by Dlamini *et al.* (2019), it was found that applying 80 tons/ha of chicken manure led to an increase in number of leaves, leaf area, leaf area index, and fresh and dry shoot mass of Swiss chard compared to lower application rates of 10, 20, and 40 t/ha. The researchers attributed the significant improvements in the parameters to the increased availability of plant nutrients in the soil, which resulted in higher manure application rates. Additionally, compost has been shown to improve soil fertility and promote beneficial microbial activity in vegetable production. Al-Kahtani *et al.* (2018) demonstrated that soil amended with a combination of compost mixture and 40% sheep manure resulted in the highest plant height, fruit yield, leaf chlorophyll, and mineral content, as well as the highest fruit quality index compared to tomato plants grown in other combinations such as 5%, 10%, and 20% sheep manure, as well as a control using active NPK mineral.

The quality characteristics of green pepper fruits, such as firmness, colour, flavour, nutritional value, and vitamin C, change during the postharvest period (El-Ramady *et al.*, 2015; Mishra and Gamage, 2020). These changes significantly influence consumer acceptance and the market value of the product. The Vitamin C content serves as a key measure of nutritional quality and antioxidant capability with higher levels linked to improved shelf life and increased health benefits for consumers. Chlorophyll levels are essential for preserving the characteristic green colour that consumers expect from fresh, high-quality peppers, and they enhance the vegetable's nutritional benefits through its antioxidant effects. Moreover, crude fibre content affects both the texture and digestibility of the peppers, playing an important role in maintaining their firmness during storage and contributing to the dietary fibre intake of consumers.

To optimise postharvest quality and storability, it is important to determine the appropriate rates of poultry manure application. This study hypothesises that different rates of poultry manure application will significantly affect the postharvest quality and storability of green pepper fruits. Specifically, moderate rates are expected to enhance physiological weight retention, shelf-life, vitamin C content, chlorophyll levels, and crude fibre content, while low or excessive rates may reduce fruit quality and increase physiological stress. The focus will be on evaluating the impact of these varied poultry manure rates on physiological weight loss, shelf-life, and nutrient composition of green pepper fruits.

MATERIALS AND METHODS

Study design

The flow chart in Figure 1 illustrates the impact of different poultry manure rates on the postharvest quality and shelf-

life of green pepper (*Capsicum annuum*) fruits. The process involved seeding, transplanting, applying poultry manure at rates of 0, 1.2, and 2.4 kg/ha, harvesting after 9 weeks, and conducting a quality analysis.

Experimental site

The experiment was conducted at the Postharvest Department of the Kumasi Institute of Tropical Agriculture in Kumasi. This site is situated at 6°40'26" North latitude and 10°35" West longitude, with an elevation of about 260 meters. The region experiences a tropical maritime climate, featuring distinct wet and dry seasons and a bimodal rainfall pattern. The primary rainy period spans March to September, peaking in June and August. November to March marks the main dry season. Annual rainfall averages 1300 mm, with a mean temperature of 28°C. The study took place during the dry season, spanning April to June 2024.

Soil and manure analysis

The soil and manure underwent analysis before cultivation. Total nitrogen was measured using the Kjeldahl method, while available phosphorus was quantified colourimetrically using a spectrophotometer. An atomic absorption spectrophotometer (AAS) determined calcium and iron levels in soil, manure, and plant tissues. Soil pH was assessed with a pH meter. All tests were performed at Crop Science Laboratory –KNUST.

Experimental design and treatments

The study employed a 3×1 factorial arrangement within a Randomized Complete Block Design (RCBD), with three replications. In all, there were 9 plots with sizes of 1.5m × 1.2m comprising an experimental field of 16.20m². Paths of 0.5m width separated both the replications and the treatments. The poultry manure treatments used include: T₁: Poultry manure (16t/ha=2.4kg); T₂: Poultry manure (8t/ha=1.2kg) and T₃: Control (0kg). The selection of application rates creates a gradient to evaluate plant responses, enabling the identification of the optimal rate for cultivating green peppers. California Wonder green pepper seeds were used for the experiment. The seeds were obtained from a Chinese-owned agrochemical shop in Kumasi. The cultivar was chosen due to its popularity among local vegetable growers and consumers. Initially, seeds were nursery-sown and after three weeks, transplanted to the field. Robust seedlings were selected and planted at 30cm × 30cm intervals, with 20 plants per plot. Regular watering and standard cultural practices were maintained throughout the growing period. The peppers were harvested at the green, fully mature stage after a total growth period of 9 weeks, including nursery time.




Figure 1. Flow chart of different poultry manure rates on the postharvest quality and shelf-life of green pepper (*Capsicum annuum*) fruits.

Sample collection and processing

After a 9-week growth period (including nursery time), the green peppers were harvested at full maturity. Six fruits (6) from each plot's centre were collected, placed in separate sterile polyethylene bags, and transported on ice to the laboratory. Quality analysis was performed at 3-day intervals (0, 3, 6, 9, and 12 days post-harvest), measuring vitamin C (mg/g), chlorophyll (mg/g), and crude fibre (%). Shelf-life evaluation was conducted over 12 days, while physiological weight loss (g) was monitored for 14 days.

Fruit quality attributes

Physiological weight loss

Physiological weight loss was measured by observing changes in the sampled fruits' weights throughout the storage period. An electronic balance was used to measure the initial and subsequent weights of pepper fruits daily. The percentage loss in weight was calculated using the method described by Haile (2018) with the following formula:

$$PLW (\%) = \frac{\text{Initial weight} - \text{Final weight}}{\text{Initial weight}} \times 100$$

Vitamin C

Vitamin C content was quantified by titrating 3g of blended pulp homogenized with 50 mL of 12% oxalic acid. The

titrant used was 2, 6-dichlorophenolindophenol sodium salt solution. Results were expressed as mg of ascorbic acid per 100g of pulp, following the method described by Abu-Zahra (2011).

Chlorophyll

Chlorophyll extraction was performed using 99.9% methanol as the solvent. Samples were stored overnight in a dark cold room at 4°C. Following extraction, leaf pigments were promptly analyzed. Absorbance was measured at 665.2, 652.4, and 470 nm. Chlorophyll content in fresh fruit was determined according to the method described by Khaleghi *et al.* (2012).

Crude fibre

Crude fibre content was determined using the methods described by Abu-Zahra *et al.* (2006), with results expressed as a percentage.

Shelf- life (days)

The produce shelf-life was determined by counting the number of days green pepper fruits were stored up to marketable or acceptable quality. The fruits were monitored for signs of wrinkling, shrinkage, spoilage and rot. The harvested fruits were placed on trays and stored at room temperature for 12 days while observing the changes critically daily.

Table 1. Soil and manure analysis.

Sample	Organic content (%)	Organic matter (%)	Total N (%)	Ca ²⁺ (cmol/k)	Mg ⁺ (Cmol/k)	Fe (mg/kg)	Available P (Mg/kg)	pH
Soil	1.596	2.75	1.04	3.0955	0.6430	16.68	2.11	6.14
Manure	14.96	25.79	2.10	8.40	2.68	11.24	2.48	8.22

Data analysis

Statistical analysis was conducted using Origin-Pro 9.2 software (Origin Lab Corporation, Northampton, MA, USA). All experiments were replicated for reliability. Results are presented as mean values with standard deviations. To identify statistically significant differences between means, Analysis of Variance (ANOVA) was performed, followed by Tukey's comparison tests. The significance level was set at 0.05%.

RESULTS AND DISCUSSION

Soil and manure analysis

Table 1 presents the pre-planting nutrient analysis of the soil and manure. Results indicate that the manure was alkaline, with a pH of 8.22, while the soil was mildly acidic with a pH of 6.14. The soil showed deficiencies in several key components: organic content, organic matter, total nitrogen, calcium, and magnesium.

Weight loss

Postharvest weight loss in fruits and vegetables is largely attributed to continuous respiration by harvested produce. The study found no significant differences ($p > 0.05$) in weight loss among green peppers from various treatments on days 0, 2, 5, 6, and 9. However, significant differences ($p < 0.05$) were observed on days 1, 3, 7, 8, 10, 11, 12, 13, and 14. By day 14, peppers from untreated plants (0 kg/ha poultry manure) showed the highest weight loss (68.7%), while those from plants treated with 2.4 kg/ha poultry manure exhibited the lowest (52.0%). No significant differences ($p > 0.05$) in weight loss were observed between peppers from plants treated with 1.2 kg/ha and 2.4 kg/ha of poultry manure. However, both these treatments resulted in significantly lower ($p < 0.05$) weight loss compared to the control group (0 kg/ha manure), as illustrated in Figure 2. The decrease in weight loss observed in peppers from plants treated with 2.4 kg/ha of fertilizer may be due to increased calcium availability. Calcium enhances fruit firmness and lowers respiration rates, which is linked to lower weight loss (Hernandez-Munoz *et al.*, 2008; Nyamah *et al.*, 2011). In a study by Angeletti *et al.* (2010), it was discovered that calcium-treated blueberries had less softening and weight loss than

control fruits. The fruits might have developed thicker pericarps, enhancing protection against moisture loss and physical damage, thus minimizing storage weight loss. Conversely, Addo *et al.* (2021) suggested that high storage weight loss percentages may result from low biomass accumulation.

Vitamin C

Vitamin C, an essential nutrient and powerful antioxidant in both plants and humans, was analyzed in stored pepper fruits. No significant differences ($p > 0.05$) in vitamin C content were observed on day 6 of storage. However, significant variations emerged on days 0, 3, 9, and 12 across treatments. By day 12, fruits from plots treated with 2.4kg/ha poultry manure exhibited the highest vitamin C content (17.89 mg/g), while those from untreated plots had the lowest (9.11 mg/g). Interestingly, there were no significant differences ($p > 0.05$) between the control and other treatments at day 12. The observation that green peppers treated with 2.4 kg/ha of poultry manure had the highest vitamin C content after harvest and storage suggests that this application rate, combined with the prevailing light and temperature conditions, created an optimal environment for nutrient uptake and vitamin C synthesis. (Figure 3). This finding aligns with Ansa and Woke's (2018) study, which reported higher vitamin C content in cayenne peppers treated with 20 t/ha manure compared to lower rates or no treatment. They concluded that increased poultry manure application correlates with higher vitamin C concentrations. Shafeek *et al.* (2015) noted that optimal organic fertilizer rates can boost fruit vitamin C accumulation, while insufficient rates may lead to nutritional deficiencies, hindering plant growth and vitamin C synthesis. A gradual decrease in ascorbic acid content was observed over the storage period, potentially due to exposure to excessive light, temperature fluctuations, physical damage, and low relative humidity, as claimed by Asanda *et al.* (2017) and Lee and Kader (2000).

Chlorophyll

The chlorophyll content of green pepper plants treated with different manure rates showed no significant differences ($p > 0.05$) at day 0 and 12. However, the results showed significant differences ($p < 0.05$) among treatments on




Figure 2. Effect of poultry manure rates on weight loss of green pepper fruits after harvest and during storage.




Figure 3. Effect of poultry manure rates on the Vitamin C content in green pepper fruits after harvest and during storage.

chlorophyll content in the pepper fruits at days 3, 6 and 9. The highest initial value (0.246 mg/g) of chlorophyll content after harvest was obtained from fruits treated with 1.2 kg/ha of poultry manure and the lowest from plots treated with no manure (0.111 mg/g). After storage, fruits from plots amended with 1.2 kg/ha manure retained the highest chlorophyll content (0.103 mg/g) while fruits treated with no manure had the lowest chlorophyll content (0.076 mg/g) (Figure 4). The higher chlorophyll content in fruits treated with 1.2 kg/ha of poultry manure may be attributed to a balanced nutrient supply, particularly

nitrogen, in the manure. Excessive nutrients can lead to toxicity and reduced chlorophyll synthesis. Moreover, the fruits appeared to regulate their physiological processes better under varying environmental conditions such as temperature, light intensity, and water availability, leading to higher chlorophyll content throughout the storage period. However, these findings contradict the results of Okonwu and Mensah (2011) who suggested applying poultry manure at a higher rate of 400 kg/ha. They observed that this higher rate resulted in high chlorophyll content, influenced by nitrogen levels.




Figure 4. Effect of poultry manure rates on the chlorophyll content in green pepper fruits after harvest and during storage.




Figure 5. Effect of poultry manure rates on the crude fibre content in green pepper fruits after harvest and during storage.

Crude fibre

The crude fibre content was not significantly ($p>0.05$) affected by the different manure rates at day 0, 3 and 6. However, the results showed significant ($p<0.05$) differences in crude fibre contents at days 9 and 12. At day 12, fruits from plots treated with 2.4 kg/ha poultry manure had the highest crude fibre content (2.09 %) while those treated with 0 kg/ha poultry manure had the lowest crude fibre content (1.21 %) at the end of storage (Figure 5).

There were no significant ($p>0.05$) differences in the crude fibre content produced by green pepper treated with 1.2 kg/ha and 2.4 kg/ha poultry manure although at day 12, both were significantly ($p<0.05$) higher than fruits treated with 0 kg/ha manure. The higher initial and final values of crude fibre contents recorded in pepper fruits treated with poultry manure rate of 2.4 kg/ha may be attributable to the increased availability of nitrogen in the manure stimulating cell division and elongation, resulting in higher crude fibre content in the pepper fruits (Taiz and Zeiger, 2022). In a

Table 2. Effects of poultry manure rates on shelf-life of green pepper fruits at storage.

Treatments	Shelf-life (days)
0 kg/ha pm	7.0 ^a
1.2 kg/ha pm	8.0 ^{ab}
2.4 kg/ha pm	10.0 ^b

study by Rehman *et al.* (2020) and Khan *et al.* (2017), it was revealed that nitrogen is crucial for the synthesis of amino acids, proteins, and chlorophyll. These compounds contribute to increased biomass and fibre production in plants.

Shelf-life

The shelf-life of green peppers refers to the length of time the pepper fruits can be stored while maintaining a marketable or acceptable quality. The storage of green pepper fruits was significantly ($p \leq 0.05$) affected by the different application rates of poultry manure. The longest storage life of 10 days was observed in fruits from plots amended with 2.4 kg/ha of poultry manure, whereas fruits from plots with no amendment (control) had the shortest storage life of 7 days (Table 2). The peppers began wilting and shrivelling on the 4th and 5th days after storage with signs of darkening, browning, and rotteness, which marked the end of the shelf-life of the fruits being observed on days 7, 8, and 10 for the 0 (control), 1.2 and 2.4 kg/ha poultry manure rates, respectively. The ability of 2.4 kg/ha poultry manure fields to produce fruits with the longest shelf life may be due to relatively high and readily available calcium levels in the manure rate having the propensity to extend fruit shelf-life. Santhosh *et al.* (2022) reported that calcium (Ca) has the potential to reduce fruit softening and postharvest decay, delay fruit ripening, delay fruit senescence, and extend the storage life of fruits. Similarly, longer storage life can be adduced to lower respiration rates by the produce since high respiration activity leads to loss of food reserves, loss of water and produce senescence (Mangaraj and Goswami, 2009; Nayik and Muzaffar, 2014). The extended shelf life of green peppers, resulting from 2.4 kg/ha poultry manure application, offers significant economic benefits for farmers and retailers. For farmers, a longer shelf life means reduced postharvest losses. Green peppers can maintain marketable quality for up to 10 days instead of just 7, allowing farmers to take advantage of a wider market window. This enables better timing of sales, potentially leading to higher prices as they can sell when demand is high, rather than being forced to sell quickly to avoid spoilage.

Conclusion

This research elucidates the significant impact of poultry manure application on the postharvest attributes of green

pepper (*Capsicum annuum* L.) fruits. The 2.4 kg/ha treatment consistently outperformed other treatments in key postharvest parameters, including weight retention, storage longevity, and nutrient content (vitamin C and crude fibre). However, the 1.2 kg/ha treatment yielded the highest chlorophyll content post-storage. Based on these results, applying 2.4 kg/ha of poultry manure was recommended since it enhances the quality of green peppers after harvest and extends their storage life. This study has important implications for sustainable farming and the use of organic fertilizers. The research shows that specific levels of poultry manure can enhance the nutritional and storage qualities of green peppers, supporting the use of organic fertilizers in crop management for better food security and increased market value for farmers.

COMPETING INTEREST

The authors have no conflict of interest to declare

ACKNOWLEDGMENT

The authors wish to express their profound appreciation to the Postharvest Department of the Kumasi Institute of Tropical Agriculture- Kumasi and Crop Science Laboratory-KNUST for technical assistance and provision of laboratory space for the experiment.

REFERENCES

- Abu-Zahra, T. R. (2011). Influence of agricultural practices on fruit quality of bell pepper. *Pakistan Journal of Biological Sciences*, 14(18), 876.
- Abu-Zahra, T. R., Al-Ismael, K., & Shatat, F. (2006). Effect of organic and conventional systems on fruit quality of strawberry (*Fragaria x ananassa* Duch) grown under plastic house conditions in the Jordan Valley. In *International Symposium on Fresh Food Quality Standards: Better Food by Quality and Assurance*, 741, 159-171.
- Addo, I., Santo, K. G., Khalid, A. A., & Ackah, E. (2021). Effects of different soil amendments and age of transplant on the post-harvest quality and shelf life of sweet pepper (*Capsicum annuum*) fruits. *Bulletin of the National Research Centre*, 45, 1-8.
- Akinfasoye, J. A. (2019). Growth, yield and nutritional quality of long cayenne pepper (*Capsicum frutescens* L.) as influenced by organic fertilizers in Ibadan and Ogbomoso, Nigeria (Doctoral dissertation, Department of Agronomy, University of Ibadan).
- Al-Kahtani, S. H., Ahmed, M. A., Al-Selwey, W. A., & Abdel-Razzak, H. S. (2018). Evaluation of composted agricultural crop wastes application on growth, mineral content, yield, and fruit quality of tomato. *Journal of Experimental Biology and Agricultural Sciences*, 6(1), 159-167.
- Anaya-Esparza, L. M., Mora, Z. V.-d.I., Vázquez-Paulino, O., Ascencio, F., & Villarruel-López, A. (2021). Bell peppers (*Capsicum annuum* L.) losses and wastes: Source for food and

- pharmaceutical applications. *Molecules*, 26, 5341.
- Angeletti, P., Castagnasso, H., Miceli, E., Terminiello, L., Concellón, A., Chaves, A., & Vicente, A. R. (2010). Effect of preharvest calcium applications on postharvest quality, softening and cell wall degradation of two blueberry (*Vaccinium corymbosum*) varieties. *Postharvest Biology and Technology*, 58(2), 98-103.
- Ansa, J. E. O., & Woke, C. (2018). Effect of spacing and poultry manure rates on growth, yield and quality of cayenne pepper (*Capsicum frutescens* L.) in Southern Rain Forest of Nigeria. *International Journal of Environment, Agriculture and Biotechnology*, 3, 1234.
- Arshad, M. J., Khan, M. I., & Ali, M. H. (2024). Enhanced wheat productivity in saline soil through the combined application of poultry manure and beneficial microbes. *BMC Plant Biology*, 24, 423.
- Asanda, M., Lembe, S. M., Samson, Z. T., & Umezuruike, L. O. (2017). Postharvest factors affecting vitamin C content of citrus fruits: A review. *Scientia Horticulturae*, 218, 95-104.
- Brezeanu, C., Brezeanu, P. M., Stoleru, V., Irimia, L. M., Lipşa, F. D., Teliban, G. -C., Ciobanu, M. M., Murariu, F., Puiu, I., & Branca, F. (2022). Nutritional value of new sweet pepper genotypes grown in organic system. *Agriculture*, 12, 1863.
- Cairo-Cairo, P., Diaz-Martin, B., Machado-de-Armas, J., & Rodriguez-Lopez, O. (2023). Effects of poultry manure on structure and some indicators of fertility in tropical soils. *Archives of Agronomy and Soil Science*, 69(13), 2692-2702.
- Danfullah, T. B., Joseph, A., Abdullahi, I.A., Victoria, M.Y., Yakubu, M., Bala, A., Ahmed, I. G., & Nwankwo, C. T. (2022). Effect of induced variation on nutritional composition of *Capsicum annum* (L.). *Journal of Science and Technology*, 6 (2).
- Dlamini, C., Masarirambi, M. T., Wahome, P. K., & Oseni, T. O. (2020). The Effects of Chicken Manure Application Rates on Growth and Yield of Swiss Chard (*Beta vulgaris* var. cicla L.). *Asian Journal of Advances in Agricultural Research*, 12(4), 12-19.
- Drózd, D., Wystalska, K., Malińska, K., Grosser, A., Grobelak, A., and Kacprzak, M. (2020). Management of poultry manure in Poland—Current state and future perspectives. *Journal of Environmental Management*, 264, 110327.
- El-Ramady, H. R., Domokos-Szabolcsy, É., Abdalla, N. A., Taha, H. S., & Fári, M. (2015). Postharvest management of fruits and vegetables storage. *Sustainable Agriculture Reviews*, 15, 65-152.
- Hernandez-Munoz, P., Almenar, E., Del Valle, V., Velez, D., & Gavara, R. (2008). Effect of chitosan coating combined with postharvest calcium treatment on strawberry (*Fragaria x ananassa*) quality during refrigerated storage. *Food Chemistry*, 110(2), 428-435.
- Jamshidovich, A. S. (2023). Ascorbic Acid: Its role in immune system, chronic inflammation diseases and on the antioxidant effects. *European Journal of Modern Medicine and Practice*, 3(11), 57-60.
- Khaleghi, E., Arzani, K., Moallemi, N., & Barzegar, M. (2012). Evaluation of chlorophyll content and chlorophyll fluorescence parameters and relationships between chlorophyll a, b and chlorophyll content index under water stress in *Olea europaea* cv. Dezful. *International Journal of Agricultural and Biosystems Engineering*, 6(8), 636-639.
- Khan, A., Tan, D. K. Y., Afridi, M. Z., Luo, H., Tung, S. A., Ajab, M. and Fahad, S. (2017). Nitrogen fertility and abiotic stresses management in cotton crop: a review. *Environmental Science and Pollution Research*, 24, 14551-14566.
- Lee, S. K., & Kader, A. A. (2000). Preharvest and postharvest factors influencing vitamin C content of horticultural crops. *Postharvest Biology and Technology*, 20(3), 207-220.
- Mangaraj, S., & Goswami, T. K. (2009). Modified atmosphere packaging of fruits and vegetables for extending shelf-life- A review. *Fresh produce*, 3(1), 1-31.
- Mishra, V. K., & Gamage, T. V. (2020). *Postharvest physiology of fruits and vegetables*, 3rd edition. Imprint CRC. p.20.
- Nayik, G. A., & Muzaffar, K. (2014). Developments in packaging of fresh fruits-shelf life perspective: A review. *American Journal of Food Science and Nutrition Research*, 1(5), 34-39.
- Ngongolo, N., & Mrimi, D. (2024). Optimizing organic agriculture: Poultry manure dominates in enhancing germination and growth of common beans in Tanzania's semi-arid environment. *Berkala Penelitian Hayati*, 30(1), 35-39.
- Nyamah E. Y., Maalekuu, B. K., & Oppong-Sekyere, D. (2011). Influence of different soil amendments on postharvest performance of tomato (*Lycopersicon esculentum*). *Journal of Stored Products and Postharvest Research*, 3(1), 11-13.
- Okonwu, K. and Mensah, S. I. (2011). Effects of poultry manure on nutrient uptake of *Cucurbita moschata* (Duch. ex Lam.) Duch. ex Poir. *Scientia Africana*, 10 (2), 16-23.
- Rehman, M., Yang, M., Fahad, S., Saleem, M. H., Liu, L., Liu, F., & Deng, G. (2020). Morpho-physiological traits, antioxidant capacity, and nitrogen metabolism in ramie under nitrogen fertilizer. *Agronomy Journal*, 112 (4), 2988-2997.
- Risse, L. M., Cabrera, M. L., Franzluebbers, A. J., Gaskin, J. W., Gilley, John, E., Killorn, R., Radcliffe, D. E., Tollner, W. E. and Zhang, H. (2006)". Land Application of Manure for Beneficial Reuse". *Biological Systems Engineering: Papers and Publications*, 65, 283-316.
- Santhosh, S., Chitdeshwari, T., & Chinnasamy, K. (2022). Calcium influences biochemical and antioxidant enzymatic activities in tomato fruits during storage. *Journal of Applied Biology and Biotechnology*, 1-7.
- Shafeek, M. R., Shaheen, A. M., Abd El-Samad, E. H., Rizk, F. A., & Abd El-Al, F. S. (2015). Response of growth, yield and fruit quality of cantaloupe plants (*Cucumis melo* L.) to organic and mineral fertilization. *Middle East Journal of Applied Sciences*, 5(1), 76-82.
- Taiz, L., & Zeiger, E. (2002). *Plant physiology*. Sinauer Associates Inc Publishers Sunderland, Massachusetts, Pp. 1-690.